

Mixed-Mean Inequality for Submatrix

Lin Si, Suyun Zhao

Abstract. For a $m \times n$ matrix $B = (b_{ij})_{m \times n}$ with nonnegative entries b_{ij} and any $k \times l$ -submatrix B_{ij} of B , let $a_{B_{ij}}$ and $g_{B_{ij}}$ denote the arithmetic mean and geometric mean of elements of B_{ij} respectively. It is proved that if k is an integer in $(\frac{m}{2}, m]$ and l is an integer in $(\frac{n}{2}, n]$ respectively, then

$$\left(\prod_{\substack{i=k, j=l \\ B_{ij} \subset B}} a_{B_{ij}} \right)^{\frac{1}{C_m^k \cdot C_n^l}} \geq \frac{1}{C_m^k \cdot C_n^l} \left(\sum_{\substack{i=k, j=l \\ B_{ij} \subset B}} g_{B_{ij}} \right),$$

with equality if and only if b_{ij} is a constant for every i, j .

1. Introduction

Let $x_1, \dots, x_n$ be positive real numbers, then the arithmetic-geometric mean inequality is

$$\frac{x_1 + \dots + x_n}{n} \geq \sqrt[n]{x_1 \cdots x_n},$$

with equality if and only if $x_1 = \dots = x_n$.

There are many research articles devoted to the classical arithmetic-geometric mean inequality and its generalizations([1],[2],[4],[9],[10]). The mixed-type arithmetic-geometric mean inequalities([3],[5],[6],[7],[8],[11]) are one of the most important branch in these generalizations.

In [6], Kedlaya established the following mixed mean inequality, which was conjectured by F.Holland and was given an inductive proof by T.Matsuda[8]:

The arithmetic mean of the numbers

$$x_1, \sqrt{x_1 x_2}, \sqrt[3]{x_1 x_2 x_3}, \dots, \sqrt[n]{x_1 x_2 \cdots x_n}$$

does not exceed the geometric mean of the numbers

$$x_1, \frac{x_1 + x_2}{2}, \frac{x_1 + x_2 + x_3}{3}, \frac{x_1 + x_2 + \dots + x_n}{n},$$

with equality if and only if $x_1 = \dots = x_n$.

2000 Mathematics Subject Classification: 26B25.

Key words and Phrases: Mixed mean, power mean, matrix, arithmetic-geometric mean inequality.

In [3], Carlson established the following mixed mean inequality:

Let the arithmetic and geometric means of the real nonnegative numbers $x_1, \dots, x_n$ taken $n - 1$ at a time be denoted by

$$a_i = \frac{x_1 + \dots + x_n - x_i}{n - 1}, \quad g_i = \left(\frac{x_1 \cdots x_n}{x_i} \right)^{\frac{1}{n-1}}.$$

Then for $n \geq 3$,

$$\left(a_1 \cdots a_n \right)^{\frac{1}{n}} \geq \frac{g_1 + \dots + g_n}{n},$$

with equality if and only if $x_1 = \dots = x_n$.

In [7], Leng, Si and Zhu generalized the above result to any subsets and established another mixed arithmetic-geometric mean inequality:

Let $X = \{x_1, \dots, x_n \mid x_i > 0, i = 1, 2, \dots, n\}$. For $A \subset X$, let a_A and g_A denote the arithmetic mean and geometric mean of all elements of A , respectively. If k is an integer in $(\frac{n}{2}, n]$, then

$$\left(\prod_{\substack{|A|=k \\ A \subset X}} a_A \right)^{\frac{1}{C_n^k}} \geq \frac{1}{C_n^k} \left(\sum_{\substack{|A|=k \\ A \subset X}} g_A \right), \quad (1)$$

with equality if and only if $x_1 = \dots = x_n$.

In this note, we established a new mixed arithmetic-geometric mean inequality for submatrix, which was an extension of the Carlson inequality and also an extension of (1).

Our main result is the following theorem.

Theorem For a $m \times n$ matrix $B = (b_{ij})_{m \times n}$ with nonnegative entries $b_{ij} \geq 0$ and any $k \times l$ -submatrix B_{ij} of B , let $a_{B_{ij}}$ and $g_{B_{ij}}$ denote the arithmetic mean and geometric mean of elements of B_{ij} respectively. If k is an integer in $(\frac{m}{2}, m]$ and l is an integer in $(\frac{n}{2}, n]$ respectively, then

$$\left(\prod_{\substack{i=k, j=l \\ B_{ij} \subset B}} a_{B_{ij}} \right)^{\frac{1}{C_m^k \cdot C_n^l}} \geq \frac{1}{C_m^k \cdot C_n^l} \left(\sum_{\substack{i=k, j=l \\ B_{ij} \subset B}} g_{B_{ij}} \right), \quad (2)$$

with equality if and only if b_{ij} is a constant for every i, j .

Remark 1. If $k \leq [\frac{m}{2}]$, for the matrix $B = (b_{ij})_{m \times n}$, taking $b_{11} = b_{12} = \dots = b_{1n} = b_{21} = b_{22} = \dots = b_{2n} = \dots = b_{k1} = b_{k2} = \dots = b_{kn} = 1$, $b_{ij} = 0$ for $k+1 \leq i \leq m$, then the right-hand side of (2) equals $1/(C_m^k \cdot C_n^l)$, but the left-hand side is zero. If $l \leq [\frac{n}{2}]$, by the same argument, one can get a contradiction. Hence the statement of Theorem fails for $k \leq [\frac{m}{2}]$ or $l \leq [\frac{n}{2}]$.

Remark 2. Taking $k = m, l = n$ in Theorem, inequality (2) is just the classical arithmetic-geometric mean inequality. For $m = 1$ or $n = 1$ in Theorem, inequality (2) is just the inequality (1).

Remark 3. The condition of our theorem is weaker than the ones of (1), because the infimum of $k \times l$ is $\frac{m}{2} \times \frac{n}{2}$, which is less than $\frac{1}{2} \times m \times n$, the half of the element number of the matrix $B = (b_{ij})_{m \times n}$.

2. Proof of Main Results

Let X denote the finite set with positive real numbers $x_1, x_2, \dots, x_n$ and let X_i denote the subset of X with k elements $x_{i_1}, \dots, x_{i_k}$. The r power mean of the elements of X_i is denoted by

$$m_r(X_i) = \left[\frac{1}{k} (x_{i_1}^r + \dots + x_{i_k}^r) \right]^{\frac{1}{r}}, \quad (3)$$

where $r > 0$ is some real number. If $r = 1$ in the above equality, we get a_{X_i} , the arithmetic mean of the elements of X_i . Let $r \rightarrow 0$ in (3), we get g_{X_i} , the geometric mean of the elements of X_i .

We first established the following lemma.

Lemma For any $m \times n$ matrix $X = (x_{ij})_{m \times n}$ with nonnegative entries $x_{ij} \geq 0$. Denote by $X_1, \dots, X_{C_m^k \cdot C_n^l}$ its all $k \times l$ -submatrix X_{ij} of X . If k is an integer in $(\frac{m}{2}, m]$ and l is an integer in $(\frac{n}{2}, n]$ respectively, then

$$m_r(X_i) = \left[\frac{1}{C_m^k \cdot C_n^l} [(m_r(X_i \cap X_1))^r + (m_r(X_i \cap X_2))^r + \dots + (m_r(X_i \cap X_{C_m^k \cdot C_n^l}))^r] \right]^{\frac{1}{r}}. \quad (4)$$

Here the intersection of $X_i \cap X_j, j = 1, \dots, C_m^k \cdot C_n^l$, is just the general set intersection.

Proof. It suffices to prove the equality

$$(m_r(X_i))^r = \frac{1}{C_m^k \cdot C_n^l} [(m_r(X_i \cap X_1))^r + (m_r(X_i \cap X_2))^r + \dots + (m_r(X_i \cap X_{C_m^k \cdot C_n^l}))^r]. \quad (5)$$

Assume that $X_i = \{x_{i_1}, \dots, x_{i_{k \times l}}\}$, then the left hand of (5) is that

$$(m_r(X_i))^r = \frac{1}{k \times l} (x_{i_1}^r + \dots + x_{i_{k \times l}}^r).$$

Now we need show that the right hand of (5) is also the mean of $\{x_{i_1}^r, \dots, x_{i_{k \times l}}^r\}$ with the same coefficient $\frac{1}{k \times l}$.

Assume that the right hand of (5) is $c_{i_1} x_{i_1}^r + \dots + c_{i_{k \times l}} x_{i_{k \times l}}^r$, by the arbitrariness of X_i , we get that $c_{i_1} = \dots = c_{i_{k \times l}}$.

Since $k > \frac{m}{2}$ and $l > \frac{n}{2}$, we have

$$X_i \cap X_j \neq \Phi, \quad j = 1, 2, \dots, C_m^k \cdot C_n^l.$$

Then the sum of all coefficients in $(m_r(X_i \cap X_j))^r$ is 1, $j = 1, 2, \dots, C_m^k \cdot C_n^l$. As a result, the sum of all coefficients in the right hand of (5) is

$$\frac{1}{C_m^k \cdot C_n^l} \underbrace{(1 + 1 + \dots + 1)}_{C_m^k \cdot C_n^l} = 1,$$

i.e., $c_{i_1} + \dots + c_{i_{k \times l}} = 1$. Therefore $c_{i_1} = \dots = c_{i_{k \times l}} = \frac{1}{k \times l}$. $\square$

Remark 1. (4) is the generalization of related result in [7]. But the condition in Lemma is weaker, because the infimum of $k \times l$ is $\frac{m}{2} \times \frac{n}{2}$, which is less than $\frac{1}{2} \times m \times n$, the half of the element number of the matrix $X = (x_{ij})_{m \times n}$.

Remark 2. In (4), if $r = 1$, we have

$$a_{X_i} = \frac{1}{C_m^k \cdot C_n^l} (a_{X_i \cap X_1} + a_{X_i \cap X_2} + \dots + a_{X_i \cap X_{C_m^k \cdot C_n^l}}). \quad (6)$$

In (4), if $r \rightarrow 0$, we have

$$g_{X_i} = (g_{X_i \cap X_1} \cdot g_{X_i \cap X_2} \cdot \dots \cdot g_{X_i \cap X_{C_m^k \cdot C_n^l}})^{\frac{1}{C_m^k \cdot C_n^l}}. \quad (7)$$

Proof of Theorem.

For a $m \times n$ matrix $B = (b_{ij})_{m \times n}$ with nonnegative entries $b_{ij} \geq 0$ and any $k \times l$ -submatrix B_{ij} of B , which is composed by the lines $(i_1, i_2, \dots, i_k)$ of B and by the rows $(j_1, j_2, \dots, j_l)$ of B , by the partial order of $(i_1, i_2, \dots, i_k)$ and $(j_1, j_2, \dots, j_l)$, we get a subset chain $B_1, B_2, \dots, B_{C_m^k \cdot C_n^l}$

In Lemma, let X be B and $X_i, j = 1, 2, \dots, C_m^k \cdot C_n^l$, be corresponding to $B_1, B_2, \dots, B_{C_m^k \cdot C_n^l}$, then (6) is

$$a_{B_i} = \frac{1}{C_m^k \cdot C_n^l} (a_{B_i \cap B_1} + a_{B_i \cap B_2} + \dots + a_{B_i \cap B_{C_m^k \cdot C_n^l}}), \quad (8)$$

and (7) is

$$g_{B_i} = (g_{B_i \cap B_1} \cdot g_{B_i \cap B_2} \cdot \dots \cdot g_{B_i \cap B_{C_m^k \cdot C_n^l}})^{\frac{1}{C_m^k \cdot C_n^l}}. \quad (9)$$

By the arithmetic-geometric inequality, it follows that

$$a_{B_i \cap B_j} \geq g_{B_i \cap B_j}, \quad j = 1, 2, \dots, C_m^k \cdot C_n^l. \quad (10)$$

From (8) and (10), we infer that

$$a_{B_i} \geq \frac{1}{C_m^k \cdot C_n^l} (g_{B_i \cap B_1} + g_{B_i \cap B_2} + \dots + g_{B_i \cap B_{C_m^k \cdot C_n^l}}).$$

Therefore

$$\begin{aligned}
\left(\prod_{\substack{i=k, j=l \\ B_{ij} \subset B}} a_{B_{ij}} \right)^{\frac{1}{C_m^k \cdot C_n^l}} &= \left(\prod_{i=1}^{C_m^k \cdot C_n^l} a_{B_i} \right)^{\frac{1}{C_m^k \cdot C_n^l}} \\
&\geq \frac{1}{C_m^k \cdot C_n^l} \left(\prod_{i=1}^{C_m^k \cdot C_n^l} \left(\sum_{j=1}^{C_m^k \cdot C_n^l} g_{B_i \cap B_j} \right) \right)^{\frac{1}{C_m^k \cdot C_n^l}}.
\end{aligned} \tag{11}$$

On the other hand, using the discrete case of Hölder's inequality in the form

$$\sum_{k=1}^n \left(\prod_{j=1}^m x_{jk} \right)^{\frac{1}{m}} \leq \left(\prod_{j=1}^n \left(\sum_{k=1}^m x_{jk} \right) \right)^{\frac{1}{m}},$$

where n, m are positive integers and $x_{jk} > 0 (j, k = 1, 2, \dots, m)$, we obtain

$$\left(\prod_{i=1}^{C_m^k \cdot C_n^l} \left(\sum_{j=1}^{C_m^k \cdot C_n^l} g_{B_i \cap B_j} \right) \right)^{\frac{1}{C_m^k \cdot C_n^l}} \geq \sum_{i=1}^{C_m^k \cdot C_n^l} \left(\prod_{j=1}^{C_m^k \cdot C_n^l} g_{B_i \cap B_j} \right)^{\frac{1}{C_m^k \cdot C_n^l}}. \tag{12}$$

Combining (9), (11) and (12), it follows that

$$\begin{aligned}
\left(\prod_{\substack{i=k, j=l \\ B_{ij} \subset B}} a_{B_{ij}} \right)^{\frac{1}{C_m^k \cdot C_n^l}} &\geq \frac{1}{C_m^k \cdot C_n^l} \sum_{i=1}^{C_m^k \cdot C_n^l} \left(\prod_{j=1}^{C_m^k \cdot C_n^l} g_{B_i \cap B_j} \right)^{\frac{1}{C_m^k \cdot C_n^l}} \\
&= \frac{1}{C_m^k \cdot C_n^l} \sum_{i=1}^{C_m^k \cdot C_n^l} g_{B_i} = \frac{1}{C_m^k \cdot C_n^l} \left(\sum_{\substack{i=k, j=l \\ B_{ij} \subset B}} g_{B_{ij}} \right)
\end{aligned}$$

which is just the inequality (2). □

References

- [1] E.F.Beckenbach and R.Bellman, *Inequalities*, Springer, Berlin, 1961.
- [2] P.S.Bullen, *A dictionary of inequalities*, Longman, 1998.
- [3] B.C.Carlson, R.K.Meany, S.A.Nelson, *An inequality of mixed arithmetic and geometric means*, SIAM Review **13**(2), 1971, 253-255.
- [4] G.H.Hardy, J.E.Littlewood and G.Pólya, *Inequalities*, second edition, Cambridge University Press, Cambridge, 1952.

- [5] Y.J.Hu, X.P.Zhang, Z.H.Yang, *Mixed mean inequalities for several positive definite matrices*, Linear Algebra and its Applications **395**(2005), 247-263.
- [6] K. Kedlaya, *Proof of a mixed arithmetic-mean, geometric-mean inequality*, Amer. Math. Monthly **101**(1994), 355-357.
- [7] G.S.Leng, L.Si, Q.S.Zhu, *Mixed-mean inequalities for subsets*, Proc. Amer. Math. Soc. **132**(2004), 2655-2660.
- [8] T. Matsuda, *An inductive proof of a mixed arithmetic-geometric mean inequality*, Amer.Math. Monthly **102**(1995), 634-637.
- [9] D.S.Mitrinović, *Analytic inequalities*, Springer, New York, 1970.
- [10] D.S.Mitrinović, J.E.Pečarić and A.M.Fink, *Classical and new inequalities in analysis*, Kluwer, Dordrecht, 1993.
- [11] B. Mond and J. Pecaric, *A mixed arithmetic-mean-harmonic-mean matrix inequality*, Linear Algebra and its Applications **237**(1996), 449-454.

Lin Si

College of Science, Beijing Forestry University,

Beijing, 100083, P.R.China

E-mail: silin@bjfu.edu.cn, lin.si@hotmail.com

Suyun Zhao

College of Mathematics & Computer Science, HeBei University,

Baoding, Hebei 071002, P.R.China

E-mail: zhaosy@mail.hbu.edu.cn